


IMAGES IN CARDIOLOGY

Pulmonary Arterial Sling Compressing the Trachea Presenting With Recurrent Stridor in an Infant

Ying Zhan, MB, Weidong Ren, MD, PhD, Yangjie Xiao, MB, Guang Song, MB, Qian Hu, MD
Shenyang, China



From the Department of
 Ultrasound, Shengjing Hospital
 of China Medical University,
 Shenyang, China.
 Manuscript received
 March 10, 2013;
 accepted March 27, 2013.

A 19-month-old female infant was referred because of recurrent stridor and was found to have a stenosed lower trachea (T) by bronchoscopy. At transthoracic echocardiography, the left pulmonary artery (LPA) was not shown in its usual location. Instead, there were 3 branches arising from the right pulmonary artery (RPA), and the left branch coursed leftward with a stenosis at the origin (A, arrow, [Online Video 1](#)). Computed tomography angiography was performed, with a confirmed diagnosis of pulmonary artery sling. The LPA was seen originating from the RPA and passing posterior to the T, leading to a compression of the origin of the LPA and the distal T (B, C, and D, arrows). These findings were directly inspected intraoperatively, and the LPA was successfully transferred to the main pulmonary artery (MPA) anterior to the T (E and F). Postoperative computed tomography angiography revealed a normal origin of the LPA and effective release from the compression (G). The infant underwent an uneventful recovery and was discharged from our hospital 1 month later. AO = aorta.